

[Current Issue Archives](#) [Cuba Health Reports](#) [MEDICC Home](#)[MEDICC Review](#) » [Spring 2008, Vol 10, No 2](#) » Research & Practice

Search MEDICC Review

Cardiovascular Risk among Older Women in a Havana Health Area

Nurys Bárbara Armas Rojas, MD, Yesenia de la Caridad Hernández Alvarez, MD, Alfredo F Dueñas Herrera, MD, Reynaldo de la Noval García, MD, Antonio Castillo Guzmán, MD, MPH

[Submit Letter to the Editors](#)[Advertise in MEDICC Review](#)[Frequently Asked Questions \(FAQs\)](#)**Information for Authors**[Manuscript Submission](#)[Author Agreement](#)[Definitions of Author & Contributor](#)[Editorial Process](#)[Peer Review](#)**Información para Autores**[Presentación de Manuscritos](#)[Convenio con el Autor](#)[Definiciones de Autor y Colaborador](#)[Proceso Editorial](#)[Arbitraje](#)**Resources for Authors**[MeSH \(Keyword Headings\)](#)[Guide for Citing References](#)[Uniform Requirements](#)[Helsinki Declaration](#)[Requisitos de uniformidad](#)**Ethics & Policies**[Editorial Ethics](#)[Open Access](#)[Ética Editorial](#)[Acceso libre](#)**About MEDICC Review**[Mission](#)[Editorial Board & Staff](#)[Peer Reviewers](#)**Introduction** Cardiovascular morbidity and mortality increase in women after menopause. Various scoring models assess qualitative risk of cardiovascular disease. The Framingham Heart Study global risk score is among the most widely used.**Objective** Determine level of coronary heart disease risk among women aged  $\geq 60$  years in a Havana health catchment area (geographic area whose residents are served by the Martires del Corynthia Polyclinic, in the Plaza de la Revolución municipality of Havana).**Methods** A descriptive, cross-sectional study was conducted in 2006. Universe: all women (3,396) aged  $\geq 60$  years in the catchment area, attended at the primary care level by the Polyclinic and 42 neighborhood family doctor-and-nurse offices. Equal probability sample: 1,082 women meeting the inclusion criteria, chosen through single-stage cluster sampling considering a  $< 10\%$  error margin for estimates for this parameter, a 95% confidence interval (CI) and a design effect of 1.5. Absolute frequencies and percentages were calculated to summarize the qualitative data obtained. Results were presented as tables.**Results** The most common cardiovascular risk factors found in this study were: physical inactivity, 74.9%; hypertension (HTN), 70.6%; abdominal obesity, 53%; reported family history of coronary heart disease (CHD), 41.8%; diabetes mellitus (DM), 21.8%; and cigarette smoking, 17.2%. Scoring according to number of risk factors present in each individual, 79.3% of these women fell into the high- or moderate-risk categories.**Conclusion** The large number of women categorized as high- or moderate-risk for coronary heart disease in this population emphasizes the need for preventive actions aimed at reducing these figures.**Keywords:** Cardiovascular diseases, vascular diseases, ischemic heart disease, coronary heart disease, postmenopause, woman, aged, risk factors, risk assessment, hypertension, high blood pressure, lifestyle, diabetes mellitus, obesity, abdominal adipose tissue, body mass index, physical inactivity, tobacco use

(Abstract)

[Download PDF](#) [See HTML](#) [HOME](#)

Article Tools

[Print this page](#)[See HTML](#)[Download PDF](#)[Like Us on Facebook](#)[Follow Us on Twitter](#)

©MEDICC, 2012

[\[Copyright\]](#) [\[Creative Commons License\]](#) [\[Privacy Policy\]](#) [\[Terms & Conditions\]](#)